

Mazuri® Waterfowl Breeder Diet

(Available at www.mazuri.com or through a Mazuri® retailer)

Formula Code 5640 | Catalog # 0001433



Description

Mazuri® Waterfowl Breeder Diet contains biologically available nutrients to enhance breeding performance, egg numbers, fecundity and chick viability for ornamental waterfowl.

Features and Benefits

- **Nutritionally balanced** – Promotes successful breeding programs and requires no mixing or supplementation.
- **Extruded floating pellet diet** - Allows for feeding of bird in the water and prevents accumulation on the bottom of the feeding area, limiting water contamination and waste and creating a more natural feeding environment.
- **Sustainable fishmeal source.**



Product Form & Packaging

Extruded feed | 3/8" diameter x 3/16" length | 50 lb. net weight paper sack

Guaranteed Analysis

Crude protein not less than	17.00%	Calcium not more than	3.50%
Crude fat not less than	2.50%	Phosphorus not less than	0.70%
Crude fiber not more than	6.00%	Salt not less than	0.05%
Ash not more than.....	11.00%	Salt not more than	0.55%
Calcium not less than.....	2.50%	Sodium not more than	0.55%

Ingredients

Ground Corn, Dehulled Soybean Meal, Wheat Middlings, Calcium Carbonate, Fish Meal, Dehydrated Alfalfa Meal, Soybean Oil, Dicalcium Phosphate, Salt, Brewers Dried Yeast, Choline Chloride, Cholecalciferol (Vitamin D3), Menadione Sodium Bisulfite Complex (Vitamin K), Vitamin A Acetate, Biotin, Manganous Oxide, d-Alpha Tocopheryl Acetate (Vitamin E), Zinc Oxide, Folic Acid, Calcium Pantothenate, Nicotinic Acid, Pyridoxine Hydrochloride, Copper Sulfate, Vitamin B12 Supplement, Riboflavin Supplement, Preserved with Mixed Tocopherols, Magnesium Oxide, Rosemary Extract, Citric Acid (a Preservative), DL-Methionine, Thiamine Mononitrate, L-Lysine, Calcium Iodate, Sodium Selenite.

Feeding Directions

Mazuri® Waterfowl Breeder Diet can be fed to birds as a major part of the daily food intake during the breeding season (spring months).

The feed can be offered ad libitum to mature birds 2 to 4 weeks before breeding season.

- Adult birds should eat in the range of 25 to 30 grams per kilogram of body weight each day, depending on species size.
- Scatter directly on the water, where it will float, or it can be fed in a long, shallow trough, which will prevent the food from floating away.
- If supplementation is desired, species-appropriate food items should be provided at no more than 20% of the total dietary intake (as is basis).

Thoroughly wash feed and water bowls on a regular basis. It is always good practice to wash hands thoroughly after feeding and/or handling animals. This diet is not for human consumption.

Important: A feeding program is only as effective as the management practices followed. Use only as directed.

Storage Conditions

For best results, store contents of open bag in container with sealing lid. Store in a cool (75°F/24°C or colder), dry (approximately 50% RH) location free from rodents and insects. Do not offer moldy or insect-infested feed to animals as it may result in performance loss, illness or death. Use within 1 year of bag manufacturing.

Notice: Repackaging of this product is prohibited and is an unauthorized use of PMI Nutrition International's trademark and product.