

# Mazuri® Waterfowl Starter Diet

(Available at [www.mazuri.com](http://www.mazuri.com) or through a Mazuri® retailer)

**Formula Code 5641 | Catalog # 0001434**



## Description

Mazuri® Waterfowl Starter Diet provides appropriate nutrition for rapid growth requirements of newly hatched chicks and adolescent birds. This diet can be fed by floating on the water or in a trough.

## Features and Benefits

- **High protein, high nutrient density diet** - Supports health and growth.
- **Special extruded form** - Floats, for a limited amount of time, for a more natural feeding.
- **Nutritionally complete** - Requires no mixing or supplementation and eliminates sorting.
- **Less waste and mess.**
- **Floating Diet** - Allows birds to be fed on the water.
- **Sustainable fishmeal source used.**



## Product Form

Extruded feed: 3/32" diameter x 3/32" length.

- 25 lb. net weight paper sack

## Guaranteed Analysis

Crude protein not less than .....	20.00%	Calcium not more than .....	1.45%
Crude fat not less than .....	3.00%	Phosphorus not less than .....	0.85%
Crude fiber not more than .....	6.50%	Salt not less than .....	0.05%
Ash not more than .....	7.00%	Salt not more than .....	0.55%
Calcium not less than .....	0.95%		

## Ingredients

Ground Corn, Wheat Middlings, Dehulled Soybean Meal, Ground Wheat, Fish Meal, Calcium Carbonate, Dicalcium Phosphate, Glyceryl Monostearate, Soybean Oil, Salt, DL-Methionine, Brewers Dried Yeast, Vitamin D3 Supplement, Pyridoxine Hydrochloride, d-Alpha Tocopheryl Acetate (Vitamin E), Menadione Sodium Bisulfite Complex (Vitamin K), Preserved with Mixed Tocopherols, Biotin, Rosemary Extract, Nicotinic Acid, Vitamin A Supplement, Choline Chloride, Manganous Oxide, Citric Acid (a Preservative), Zinc Oxide, Calcium Pantothenate, Folic Acid, Tagetes (Aztec Marigold) Extract (Color), Copper Sulfate, Thiamine Mononitrate, Riboflavin Supplement, Vitamin B12 Supplement, L-Lysine, Calcium Iodate, Sodium Selenite.

## Feeding Directions

Feed ad libitum as the sole ration to newly hatched and adolescent birds during their period of fastest growth. If supplementation is desired, it should make up no more than 20% of the total dietary intake (as-is basis).

## Storage Conditions

For best results, store contents of open bag in container with sealing lid. Store in a cool (75°F/24°C or colder), dry (approximately 50% RH) location free from rodents and insects. Do not offer moldy or insect-infested feed to animals as it may result in performance loss, illness or death. Use within 1 year of bag manufacturing.